

Antitumoral Properties of Xanthones from Mangosteen (*Garcinia Mangostana L.*) Hull

Short
ReviewCheok Choon-Yoong^{*,1}, Chin Nyuk-Ling²¹ Department of Chemical and Petroleum Engineering, Faculty of Engineering, UCSI University, Lot 12734, Jalan Choo Lip Kung, Taman Tayton View Cheras, 56000 Kuala Lumpur, Malaysia² Department of Process and Food Engineering, Faculty of Engineering, Universiti Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia**ARTICLE INFO****ABSTRACT****Article history:**

Received 7 April 2017

Received in revised form 16 May 2017

Accepted 19 May 2017

Available online 15 June 2017

Mangosteen is known as “the queen of fruits” in Malaysia, and the fruit is usually available during seasons from June to August yearly. This fruit comprises substantial amount of hull which disposes off as waste. However, it is the hull that has been discovered having various pharmaceutical properties such as antioxidant, anti-inflammatory, and antitumoral. The sudden surge of antitumoral properties studies of mangosteen hull in recent years is due to the rising of cancer which is the leading cause of death worldwide. This review highlights the recent discoveries of antitumoral properties of mangosteen hull. The major xanthones isolated from mangosteen hull which attribute to antitumoral properties are α -mangostin and γ -mangostin. They have been majorly discovered against cancers of colon, breast, and leukemia, followed by skin, bone, lung, brain, pancreatic, prostate, and head and neck.

Keywords:Mangosteen hull, Antitumoral properties, α -mangostin, γ -mangostin

Copyright © 2017 PENERBIT AKADEMI BARU - All rights reserved

1. Introduction

Mangosteen (*Garcinia mangostana L.*), is a large tropical evergreen tree from the *Guttiferae* family with a straight trunk, and it takes almost 6 to 7 years to cultivate before it fruits [1]. The fruit comprises of 25-29% of edible flesh, 60-65% of hull, and 6-10% of seed [2]. Its sweet and mild sour delicious pulp are the contributing factors for its popularity among the tropical fruits. Mangosteen has become an economically important species in recent decades since it is popular to Western Europe and United States most probably because of its pharmaceutical properties discovery, especially antitumoral [3-4]. As reported by World Health Organization, cancer is the leading cause of death worldwide and accounted for 7.6 million deaths (around 13% of all deaths) in 2008 [5]. Mangosteen is the local seasonal fruit of Malaysia, which the delicious pulp is a target by local but the deep reddish purple hull (exocarp) is disposed as waste. The hull which contains bioactive compounds of xanthones has been widely discovered having various pharmaceutical properties. Among the xanthones, α -mangostin is the most extensively discovered having inhibition capabilities against various cancer cells. This paper discusses the recent discoveries and research trend of antitumoral properties of α -mangostin on various

*Corresponding author.

E-mail address: cheokcy@ucsiuniversity.edu.my or elmayoong@gmail.com

cancer cells. The γ -mangostin and other xanthenes which have been proven as potential anticancer agents are also highlighted.

2. The α -Mangostin

The most abundant xanthone found in mangosteen hull is the α -mangostin [6-7]. α -mangostin is one of the earliest naturally occurring xanthone isolated from mangosteen hull using gas chromatography [8]. Numerous studies have revealed that the α -mangostin not only possesses properties of antioxidant, anticancer, anti-inflammatory, anti-allergy, antimicrobial, and antiparasitic, it also has potential for anti-obesity and treatment of Alzheimer's disease [9]. Within the many pharmaceutical properties, its antitumor capabilities against various cancer cells have been extensively researched in recent decades. **Fig. 1** illustrates recent scientific findings of antitumoral properties of α -mangostin against various cancer cells. Approximately 25% of previous studies have proven the inhibition capabilities of α -mangostin against colon cancer cells as shown in **Fig. 1**. Its inhibition effects on other cancer cells such as breast cancer, leukemia, skin cancer, prostate, bone, lung, brain, pancreatic, head and neck have also been investigated and discovered.




Fig. 1. Recent discoveries of antitumoral properties of α -mangostin on various cancers

Fig. 2 demonstrates the trend of antitumoral properties discovery of α -mangostin from 2000 to 2014. The earliest finding on antitumoral properties of α -mangostin were on leukemia [10-11] and colon [12-13] cell lines from 2003 to 2005. For the period from 2006 to 2010, α -mangostin's inhibition effects against breast cancer [14-16], prostate [17], and lung [18] was reported. Most recently from year 2011 to 2014, the finding of α -mangostin's inhibition for cancer cells have extended to bone [19], brain [20], skin [21-23], pancreatic [24], head and neck [25], apart from breast cancer [26-27]. It is noteworthy that α -mangostin has been extensively discovered its inhibition ability against colon cancer cell lines [28-29] for a decade.




Fig. 2: Research trend of antitumoral properties of α -mangostin derived from mangosteen hull

3. The γ -Mangostin and other Xanthenes

The γ -mangostin is another xanthone derived from mangosteen hull and for its widely studied pharmaceutical property. It appears as a paleyellow powder after isolation with a thin layer chromatography [30]. The γ -mangostin isolated from mangosteen hull has been discovered to have antitumoral property against leukemia [10, 31], breast cancer [14], skin [21], and colon cancer [28, 32].

Further to this, the 1,3,6,7-tetrahydroxy-2,8-(3-methyl-2-butenyl) xanthone and epicatechin that were studied and tested *in vitro* also showed good cytotoxicities against human breast and colon cancer cells [33]. Both have been suggested as potential anticancer agents [33]. Other two newly identified xanthenes, the tetrahydroxanthone and garcimangosxanthone were revealed to exhibit *in vitro* cytotoxicity against human lung cancer, pulmonary carcinoma, and hepatoma [34]. More recently, xanthone of 8-deoxygartanin derived from mangosteen pericarp has been discovered having cytotoxicity against skin cancer [21].

4. Conclusion

Cancer, as the leading cause of death worldwide in recent decades, has triggered the quest of plant materials which contain potential antitumoral properties to combat the disease. The α -mangostin derived from mangosteen has natural occurring bioactive compounds which is anti-cancer. The recent research trend on the discovery of this compound provides an insight for scientists or researchers to further explore its inhibition capabilities against more cancer cell lines.

References

- [1] Osman, M. and Milan, A.R. Taxonomy. In: Mangosteengarciniamangostana L., ed. J.T. Williams et al., Southampton Centre for Underutilised Crops. (2006): 1-16.
- [2] Cheok, C.Y., Mohd Adzahan, N., Abdul Rahman, R., Zainal Abedin, N.H., Hussain, N., Sulaiman, R. and Chong, G.H. "Current Trends of Tropical Fruit Waste Utilization." *Critical Reviews in Food Science and Nutrition* (2016) DOI: 10.1080/10408398.2016.1176009.
- [3] Pedraza-Chaverri, J., Cárdenas-Rodríguez, N., Orozco-Ibarra, M., Pérez-Rojas, J.M. "Medicinal properties of mangosteen (*Garcinia mangostana*)." *Food and Chemical Toxicology* 46 (2008): 3227-3239.
- [4] Ibrahim, M.Y., Hashim, N.M., Mariod A.A., Mohan, S., Abdulla, M.A., Abdelwahab, S.I. and Arbab, I.A. "α-mangostin from *Garcinia mangostana* Linn: An updated review of its pharmaceutical properties." *Arabian Journal of Chemistry* 9 (2016): 317-329.
- [5] Cancer. World Health Organization. Retrieved on 1 June 2016. <http://www.who.int/media/centre/factsheets/fs297/en/>.
- [6] Chaivisuthangkura, A., Malaikaew, Y., Chaovanalikit, A., Jaratrungrawee, A., Panseeta, P., Ratananukul, P. and Suksamrarn, S. Prenylatedxanthone composition of *Garcinia mangostana* (mangosteen) fruit hull. *Chromatographia* 69 (2008): 315-318.
- [7] Wittenauer, J., Falk, S., Schweiggert-Weisz, U. and Carle, R. "Characterisation and quantification of xanthenes from the aril and pericarp of mangosteens (*Garcinia mangostana*L.) and a mangosteen containing functional beverage by HPLC–DAD–MS". *Food Chemistry* 134 (2012): 445-452.
- [8] Jefferson, A. and Stacey, C.I. "Gas-liquid chromatography of naturally occurring xanthenes and related derivatives." *Journal of Chromatography* 57 (1971): 247-254.
- [9] Wang, Y., Xia, Z., Xu, J.R., Wang, Y.X., Hou, L.N., Qiu, Y. and Chen, H.Z. "α-Mangostin, a polyphenolic xanthone derivative from mangosteen, attenuates β-amyloid oligomers-induced neurotoxicity by inhibiting amyloid aggregation." *Neuropharmacology* 62 (2012): 871-881.
- [10] Matsumoto, K., Akao, Y., Kobayashi, E., Ohguchi, K., Ito, T., Tanaka, T., Iinuma, M. and Nozawa, Y. "Induction of apoptosis by xanthenes from mangosteen in human leukemia cell lines." *Journal of Natural Product* 66 (2003): 1124-1127.
- [11] Matsumoto, K., Akao, Y., Yi, H., Ohguchi, K., Ito, T., Tanaka, T., Kobayashi, E., Iinuma, M. and Nozawa, Y. "Preferential target is mitochondria in α-mangostin-induced apoptosis in human leukemia HL60 cells." *Bioorganic and Medicinal Chemistry* 12 (2004): 5799-5806.
- [12] Matsumoto, K., Akao, Y., Ohguchi, K., Ito, T., Tanaka, T., Iinuma, M. and Nozawa, Y. "Xanthenes induce cell-cycle arrest and apoptosis in human colon cancer DLD-1 cells." *Bioorganic & Medicinal Chemistry* 13 (2005): 6064-6069.
- [13] Nabandith, V., Suzui, M., Morioka, T., Kaneshiro, T., Kinjo, T., Matsumoto, K., Akao, Y., Iinuma, M. and Yoshimi, N. "Inhibitory effects of crude α-mangostin, a xanthone derivative, on two different categories of colon preneoplastic lesions induced by 1,2-dimethylhydrazine in the rat." *Asian Pacific Journal of Cancer Prevention* 5 (2004): 433-438.
- [14] Balunas, M.J., Su, B., Brueggemeier, R.W. and Kinghorn, A.D. "Xanthenes from the botanical dietary supplement mangosteen (*Garcinia mangostana*) with aromatase inhibitory activity." *Journal of Natural Product* 71 (2008): 1161-1166.
- [15] Lee, Y.B., Ko, K.C., Shi, M.D., Liao, Y.C., Chiang, T.A., Wu, P.F., Shih, Y.X. and Shih, Y.W. "α-Mangostin, a novel dietary xanthone, suppresses TPA-mediated MMP-2 and MMP-9 expressions through the ERK signaling pathway in MCF-7 human breast adenocarcinoma cells." *Journal of Food Science* 75 (2010): H13-H23.
- [16] Suksamrarn, S., Komutiban O., Ratananukul, P., Chimnoi, N., Lartpornmatulee, N. and Suksamrarn, A. "Cytotoxic prenylatedxanthenes from the young fruit of *Garcinia mangostana*." *Chemical and Pharmaceutical Bulletin* 54 (2006): 301-305.
- [17] Hung, S.H., Shen, K.H., Wu, C.H., Liu, C.L. and Shih, Y.W. "α-mangostin suppresses PC-3 human prostate carcinoma cell metastasis by inhibiting matrix metalloproteinase-2/9 and urokinase-plasminogen expression through the JNK signaling pathway." *Journal of Agricultural and Food Chemistry* 57 (2009): 1291-1298.
- [18] Shih, Y.W., Chien, S.T., Chen, P.S., Lee, J.H., Wu, S.H. and Yin, L.T. "α-Mangostin suppresses phorbol 12-myristate 13-acetate-induced MMP-2/MMP-9 expressions via avb3 integrin/FAK/ERK and NF-κB signalling pathway in human lung adenocarcinoma A549 cells." *Cell Biochemistry and Biophysics* 58 (2010): 31-44.

- [19] Krajarng, A., Nakamura, Y., Suksamrarn, S. and Watanapokasin, R. “ α -mangostin induces apoptosis in human chondrosarcoma cells through downregulation of ERK/JNK and Akt signaling pathway.” *Journal of Agricultural and Food Chemistry* 59 (2011): 5746-5754.
- [20] Chao, A.C., Hsu, Y.L., Liu, C.K. and Kuo, P.L. “ α -mangostin, a dietary xanthone, induces autophagic cell death by activating the AMP-activated protein kinase pathway in glioblastoma cells.” *Journal of Agricultural and Food Chemistry* 59 (2011): 2086-2096.
- [21] Wang, J.J., Sanderson, B.J.S. and Zhang, W. “Cytotoxic effect of xanthones from pericarp of the tropical fruit mangosteen (*Garcinia mangostana*Linn.) on human melanoma cells.” *Food and Chemical Toxicology* 49 (2011): 2385-2391.
- [22] Wang, J.J., Sanderson, B.J.S. and Zhang, W. “Significant anti-invasive activities of alpha-mangostin from the mangosteen pericarp on two human skin cancer cell lines.” *Anticancer Research* 32 (2012): 3805-3816.
- [23] Wang, J.J., Shi, Q.H., Zhang, W. and Sanderson, B.J.S. “Anti-skin cancer properties of phenolic-rich extract from the pericarp of mangosteen (*Garcinia mangostana*Linn.)” *Food and Chemical Toxicology* 50(2012): 3004-3013.
- [24] Hafeez, B., Mustafa, A., Fischer, J.W., Singh, A., Zhong, W., Shekhani, M.O., Meske, L., Havighurst, T., Kim, K.M. and Verma, A.K. “ α -Mangostin: A dietary antioxidant derived from the pericarp of *Garcinia mangostana* L. inhibits pancreatic tumor growth in xenograft mouse model.” *Antioxidants and Redox Signaling* 21 (2014): 682-699.
- [25] Kaomongkolgit, R., Chaisomboon, N. and Pavasant, P. “Apoptotic effect of alpha-mangostin on head and neck squamous carcinoma cells.” *Archives of Oral Biology* 56 (2011): 483-490.
- [26] Shibata M.A., Matoba, Y., Tosa, H. and Iinuma, M. “Effects of mangosteen pericarp extracts against mammary cancer.” *Alternative and Integrative Medicine* 2 (2013): 1-5.
- [27] Won, Y.S., Lee, J.H., Kwon, S.J., Kim, J.Y., Park, K.H., Lee, M.K. and Seo, K.I. “ α -Mangostin-induced apoptosis is mediated by estrogen receptor α in human breast cancer cells.” *Food and Chemical Toxicology* 66 (2014): 158-165.
- [28] Abdalrahim, F.A.A., Khalid, M.A.S., Mohammad, J.S., Zhari, I., Amin Malik, S.A.M. “*In vitro* and *in vivo* anti-colon cancer effects of *Garcinia mangostanaxanthones* extract.” *BMC Complementary and Alternative Medicine* 12 (2012): 1-10.
- [29] Nakagawa, Y., Iinuma, M., Naoe, T., Nozawa, Y., Akao, Y. “Characterized mechanism of α -mangostin-induced cell death: Caspase-independent apoptosis with release of endonuclease-G from mitochondria and increased miR-143 expression in human colorectal cancer DLD-1 cells.” *Bioorganic & Medicinal Chemistry* 15 (2007): 5620-5628.
- [30] Sen, A.K., Sarkar, K.K., Mazumder, P.C., Banerji, N., Uusvuori, R. and Hase, T.A. “The structures of garcinones a, b and c: Three new xanthones from *Garcinia mangostana*.” *Phytochemistry* 21 (1982): 1747-1750.
- [31] Itoh, T., Ohguchi, K., Iinuma, M., Nozawa, Y. and Akao, Y. “Inhibitory effect of xanthones isolated from the pericarp of *Garcinia mangostana* L. on rat basophilic leukemia RBL-2H3 cell degranulation.” *Bioorganic and Medicinal Chemistry* 16 (2008): 4500-4508.
- [32] Watanapokasin, R., Jarinthanan, F., Jerusalmi, A., Suksamrarn, S., Nakamura, Y., Sukseree, S., Uthaisang-Tanethpong-tamb, W., Ratananukul, P. and Sano, T. “Potential of xanthones from tropical fruit mangosteen as anti-cancer agents: Caspase-dependent apoptosis induction *in vitro* and in mice.” *Applied Biochemistry and Biotechnology* 162 (2010): 1080-1094.
- [33] Yu, L., Zhao, M., Yang, B. and Bai, W. “Immunomodulatory and anticancer activities of phenolics from *Garcinia mangostana* fruit pericarp.” *Food Chemistry* 116 (2009): 969-973.
- [34] Zhang, Y., Song, Z., Hao, J., Qiu, S. and Xu, Z. “Two new prenylatedxanthones and a new prenylatedtetrahydroxanthone from the pericarp of *Garcinia mangostana*.” *Fitoterapia* 81 (2010): 595-599.